

**Legend**

bedrock\_near\_surface

**gradient\_%**

- 0.060534 - 0.277187
- 0.277188 - 0.518634
- 0.518635 - 0.950187
- 0.950188 - 1.814420
- 1.814421 - 3.616536

**VF\_matLB**

- Fine
- Fine Mixed
- Coarse Mixed
- Coarse
- focus\_areas

**PMG\_dissolve**

- I >80 loess
- III 40-80 loess / till or lac
- IV 0-40 loess/till
- IX 40-80 loess (alluv?)/outwash
- Misc
- V colluv. in depressions
- VIII loamy, strat / sand
- X 0-20 alluv. / sand & gravel
- XI silt loam, lacust-alluv
- XII IL Valley outwash
- XV alluvium, silty to sandy
- XVII Misc.

Mass Wasting Observed

Center of Assessed Reaches

0 0.25 0.5 1 Miles

**Histogram Data (Estimated from PMG\_dissolve)**

| PMG_dissolve Class              | Frequency |
|---------------------------------|-----------|
| I >80 loess                     | ~10       |
| III 40-80 loess / till or lac   | ~15       |
| IV 0-40 loess/till              | ~10       |
| IX 40-80 loess (alluv?)/outwash | ~5        |
| Misc                            | ~5        |
| V colluv. in depressions        | ~5        |
| VIII loamy, strat / sand        | ~5        |
| X 0-20 alluv. / sand & gravel   | ~5        |
| XI silt loam, lacust-alluv      | ~5        |
| XII IL Valley outwash           | ~5        |
| XV alluvium, silty to sandy     | ~5        |
| XVII Misc.                      | ~5        |

